MYCOTAXON

Volume 108, pp. 49–52 *April–June 2009*

A new species of *Alternaria* on *Humulus scandens* from China

Meng Zhang[1] Hai-Yan Wu[2] & Dong-Xia Li[1]

ZM20066@126.com

[1.] *College of Plant Protection, Henan Agricultural University Zhengzhou, Henan, China, 450002*

[2.] *Center of Electronic Teaching, Henan Agricultural University Zhengzhou, Henan, China, 450002*

Abstract—A new species of the genus *Alternaria*, *A. humuli-scandens*, on *Humulus scandens* is reported. *A. humuli-scandens* is the first large-spored *Alternaria* species with long filiform rostra reported on plants of *Cannabaceae*. *A. humuli-scandens* is different from all the known species in the genus *Alternaria* based on the morphology and pathogenicity. The type specimen is deposited in the Herbarium of Henan Agricultural University: Fungi (HHAUF).

Key words—taxonomy, hyphomycetes, leaf spots, *Humulus japonicus*

In the course of a survey of mitosporic fungal pathogens of important weeds in China, a new taxon of *Alternaria* was found on *Humulus scandens* [= *Humulus japonicus* Siebold & Zucc.] in Henan Province. It is described as follows.

Alternaria humuli-scandens Meng Zhang, H.Y. Wu & Dong X. Li, **sp. nov.**

Mycobank # MB 512829 fig. 1, plate 1

Conidia solitaria, obclavata vel ovoidea, recta, laevia, pallide brunnea, septa transversa 5–8, septa longitudinalia vel obliqua 2–6, constricta, 47–75 × 14–20.5 µm, rostrum septata, hyalina, 117–195 µm longum, e basi 3–5 µm ad apicem 2 µm diam.

Holotype: HHAUF$_{06}$0529=ZM$_{06}$0529 [dried cultures on PCA, isolated from leaf spots of *Humulus scandens* (Lour.) Merr.: Zhengzhou, Henan Province, 2006, Coll. Dong X. Li and H.Y. Wu].

Leaf spots irregular, brown, 5–12 mm in diameter. Colonies on PCA grayish-brown, velvety, effuse, sporulating freely. Conidiophores arising singly, terminally or laterally from hyphae, simple, straight or curved, smooth-walled, septate, pale brown, 55–75 × 7–8 µm. Conidia solitary, obclavate or ovoid, base obtuse, straight, pale brown, smooth-walled, with 5–8 transverse septa and 2–6 longitudinal or oblique septa, constricted at some main septa, 47–75



FIGURE 1. *Alternaria humuli-scandens*
Conidia and conidiophores on PCA. (ex holotype, bar=50μm)

× 14–20.5 μm. The tapering rostra are 117–195 μm long, ca 3–5 μm wide at the base, narrowing throughout most of their length to ca 2 μm, filiform, straight, septate, hyaline.

COMMENTS: Commonly large-spored *Alternaria* species are host specific in family; host ranges can be one of the identification criteria. Up to now, only



PLATE 1 *Alternaria humuli-scandens*
Conidia and conidiophores on PCA. (ex holotype)

Alternaria humuli E.G. Simmons (a small-spored species) has been reported on plants of *Cannabaceae* (Simmons 2002, 2007). *A. humuli-scandens* is the first large-spored *Alternaria* species with long filiform rostra reported on this family. Compared to other species of *Alternaria* on the related host family *Moraceae*, *A. humuli-scandens* is close to *Alternaria fici* Farneti in size range of conidium bodies, but the conidia of the latter have beaks that are short or lacking entirely, while the conidia of *A. humuli-scandens* have long filiform rostra (Farneti 1904, Zhang et al. 2003).

Acknowledgments

The authors are grateful to Dr. E.G. Simmons, U.S.A., Prof. Y.L. Guo, Institute of Microbiology, Academia Sinica, and Prof. T.Y. Zhang, China, for reviewing the manuscript.

Literature cited

Farneti R. 1904. Le volatiche e l'atrofia dei frutti del fico. Atti. Ist. Bot. Di Pavia. 8: 513–518.

Simmons EG. 2002. *Alternaria* themes and variations (305-309) *Lewia/Alternaria* revisited. Mycotaxon 83: 127–145.

Simmons EG. 2007. *Alternaria*: an identification manual. CBS Fungal Biodiversity Centre, Utrecht.

Zhang TY, Zhang JZ, Chen WQ, Zhang M, Sun GY, Gao MX. 2003. Flora Fungorum Sinicorum, 16: *Alternaria*. BeiJing: Science Press, 1–283 + Plates (in Chinese).